

CLASS - XII

Time : Three Hours

Max. Marks : 70

General Instructions

1. All questions are compulsory.
2. Question nos. 1 to 8 are very short answer questions and carry one mark each.
3. Question nos. 9 to 18 are short answer questions and carry two marks each.
4. Question nos. 19 to 27 are also short answer questions and carry three marks each.
5. Question nos. 28 to 30 are long answer questions and carry five marks each.
6. Use log tables if necessary. Calculators are not allowed.

1. Define molality. 1
2. What are the physical states of dispersed phase and dispersion medium of froth? 1
3. Write the balanced equation for complete hydrolysis of XeF_6 . 1
4. Write the structure of :
4 - methyl pent - 3 - en - 2 - one 1
5. A compound contains two types of atoms - X and Y. It crystallises in a cubic lattice with atom X at the corners of the unit cell and atoms Y at the body centres. What is the simplest possible formula of this compound? 1
6. What is the Van't Hoff factor for a compound which undergoes tetramerization in an organic solvent? 1
7. An ore sample of galena (PbS) is contaminated with zinc blende (ZnS). Name one chemical which can be used to concentrate galena selectively by froth floatation method. 1
8. Predict the shape of ClF_3 on the basis of VSEPR theory. 1
9. Ethylene glycol (molar mass = 62 g mol^{-1}) is a common automobile antifreeze. Calculate the freezing point of a solution containing 12.4g of this substance in 100 g of water. Would it be advisable to keep this substance in the car radiator during summer?
Given : K_f for water = 1.86 K kg/mol
 K_b for water = 0.512 K kg/mol 2
10. Consider the reaction $A \rightarrow P$. The change in concentration of A with time is shown in the following plot:
(i) Predict the order of the reaction.
(ii) Derive the expression for the time required for the completion of the reaction.
11. Free energies of formation of MgO(s) and CO(g) at 1273 K and 2273K are given below
 MgO(s) = - 941 kJ/mol at 1273K
 MgO(s) = - 314 kJ/mol at 2273K
 CO(g) = - 439 kJ/mol at 1273K
 CO(g) = - 628 kJ/mol at 2273K
On the basis of above data, predict the temperature at which carbon can be used as a reducing agent for MgO(s) . 2
12. Name the two components of starch. How do they differ from each other structurally? 2
13. (a) What changes occur in the nature of egg proteins on boiling?
(b) Name the type of bonding which stabilizes α -helix structure in proteins. 2

14. Describe the mechanism of the formation of diethyl ether from ethanol in the presence of concentrated sulphuric acid. 2

15. Complete and name the following reactions:

2

16. Give chemical tests to distinguish between compounds in each of the following pairs:

(i) Phenol and Benzyl alcohol

(ii) Butane-2-ol and 2-Methyl propan-2-ol

17. Predict, giving reasons, the order of basicity of the following compounds in (i) gaseous phase and (ii) in aqueous solutions 2

OR

Account for the following:

(a) Aniline does not undergo Friedel Crafts alkylation

(b) Although -NH₂ group is an ortho and para-directing group, nitration of aniline gives alongwith ortho & para derivatives meta-derivative also.

18. Give reasons for the following :

(a) At higher altitudes, people suffer from a disease called anoxia. In this disease, they become weak and cannot think clearly.

(b) When mercuric iodide is added to an aqueous solution of KI, the freezing point is raised. 2

19. An element X with an atomic mass of 60g/mol has density of 6.23g cm⁻³. If the edge length of its cubic unit cell is

400 pm, identify the type of cubic unit cell. Calculate the radius of an atom of this element. 3

20. Write names of monomer/s of the following polymers and classify them as addition or condensation polymers.

(a) Teflon

(b) Bakelite

(c) Natural Rubber 3

21. (a) Give the IUPAC name of :

[Cr Cl₂ (H₂O)₄] Cl

(b) Give the number of unpaired electrons in the following complex ions:

(c) Name the isomerism exhibited by the following pair of coordination compounds:

Give one chemical test to distinguish between these two compounds. 3

22. Explain the following observations:

(a) Ferric hydroxide sol gets coagulated on addition of sodium chloride solution

(b) Cottrell's smoke precipitator is fitted at the mouth of the chimney used in factories.

(c) Physical adsorption is multilayered, while chemisorption is monolayered. 3

23. Account for the following:

(a) Chlorine water has both oxidizing and bleaching properties.

(b) H₃PO₂ and H₃PO₃ act as good reducing agents while H₃PO₄ does not.

(c) On addition of ozone gas to KI solution, violet vapours are obtained. 3

24. The decomposition of N₂O₅(g) is a first order reaction with a rate constant of 5 x 10⁻⁴ sec⁻¹ at 45°C. i.e. 2N₂O₅(g)

$4\text{NO}_2(\text{g}) + \text{O}_2(\text{g})$. If initial concentration of N_2O_5 is 0.25M, calculate its concentration after 2 min. Also

calculate half life for decomposition of $\text{N}_2\text{O}_5(\text{g})$.

(b) For an elementary reaction

the rate of appearance of C at time 't' is $1.3 \times 10^{-4} \text{ mol l}^{-1} \text{ s}^{-1}$.

Calculate at this time

(i) rate of the reaction.

(ii) Rate of disappearance of A. 3

25. (a) Which of the following two compounds would react faster by $\text{S}_{\text{N}}2$ path way : 1 - bromobutane or

2 - bromobutane and why.?

(b) Allyl chloride is more reactive than n - propyl chloride towards nucleophilic substitution

reaction. Explain why?

(c) Haloalkanes react with KCN to give alkyl cyanide as main product while with AgCN they form isocyanide as

main product. Give reason. 3

26. Give reasons for the following:

(a) CN^- ion is known but CP^- ion is not known.

(b) NO_2 dimerises to form N_2O_4

(c) ICl is more reactive than I_2 3

OR

An element 'A' exists as a yellow solid in standard state. It forms a volatile hydride 'B' which is a foul smelling gas

and is extensively used in qualitative analysis of salts. When treated with oxygen, 'B' forms an oxide 'C' which is

a colourless, pungent smelling gas. This gas when passed through acidified KMnO_4 solution, decolourises it. 'C'

gets oxidized to another oxide 'D' in the presence of a heterogeneous catalyst. Identify A,B,C,D, and also give the

chemical equation of reaction of 'C' with acidified KMnO_4 solution and for conversion of 'C' to 'D'.

27. Account for the following:

(a) Aspirin drug helps in the prevention of heart attack.

(b) Diabetic patients are advised to take artificial sweeteners instead of natural sweeteners.

(c) Detergents are non-biodegradable while soaps are biodegradable. 3

28. (a) An organic compound 'A' with molecular formula $\text{C}_5\text{H}_8\text{O}_2$ is reduced to n-pentane on treatment with Zn-Hg/

HCl. 'A' forms a dioxime with hydroxylamine and gives a positive Iodoform test and Tollen's test. Identify the

compound A and deduce its structure.

(b) Write the chemical equations for the following conversions:

(not more than 2 steps)

(i) Ethyl benzene to benzene

(ii) Acetaldehyde to butane - 1, 3 - diol

(iii) Acetone to propene 5

OR

(a) An organic compound 'A' with molecular formula C_8H_8O gives positive DNP and iodoform tests. It does not reduce Tollen's or fehling's reagent and does not decolourise bromine water also. On oxidation with chromic acid (H_2CrO_4), it gives a carboxylic acid (B) with molecular formula $C_7H_6O_2$. Deduce the structures of A and B.

(b) Complete the following reactions by identifying A, B and C

(i)

(ii)

29. (a) Calculate the equilibrium constant for the reaction

(b) When a current of 0.75A is passed through a $CuSO_4$ solution for 25 min, 0.369 g of copper is deposited at the

cathode. Calculate the atomic mass of copper.

(c) Tarnished silver contains Ag_2S . Can this tarnish be removed by placing tarnished silver ware in an aluminium

pan containing an inert electrolytic solution such as NaCl. The standard electrode potential for half reaction :

and for 5

OR

(a) Calculate the standard free energy change for the following reaction at $25^\circ C$

Predict whether the reaction will be spontaneous or not at $25^\circ C$. Which of the above two half cells will act as an

oxidizing agent and which one will be a reducing agent?

(b) The conductivity of 0.001M acetic acid is $4 \times 10^{-5} S / cm$. Calculate the dissociation constant of acetic acid, if

for acetic acid is $390.5 S cm^2/mol$.

30. (a) A blackish brown coloured solid 'A' when fused with alkali metal hydroxides in presence of air, produces a

dark green coloured compound 'B', which on electrolytic oxidation in alkaline medium gives a dark purple coloured

compound C. Identify A, B and C and write the reactions involved.

(b) What happens when an acidic solution of the green compound (B) is allowed to stand for some time? Give the

equation involved. What is this type of reaction called? (3 + 2 = 5)

OR

Give reasons for the following:

(a) Transition metals have high enthalpies of atomization.

(b) Among the lanthanoids, Ce(III) is easily oxidised to Ce(IV).

(c) redox couple has less positive electrode potential than couple.

(d) Copper (I) has d_{10} configuration, while copper (II) has d_9 configuration, still copper (II) is more stable in

aqueous solution than copper (I).

(e) The second and third transition series elements have almost similar atomic radii. 5

MARKING SCHEME

CHEMISTRY SAMPLE PAPER - II

CLASS - XII

Q.No. Value Points Marks

1. Number of moles of solute dissolved per K.g. of solvent (1)

2. Dispersed phase : gas (½)

Dispersion medium : liquid (½)

3. $\text{XeF}_6 + \text{H}_2\text{O} \longrightarrow \text{XeOF}_4 + \text{HF}$ (1)

4. (1)

5. XY (1)

6. (1)

7. NaCN, Sodium cyanide, used as a depressant. (1)

8. T - shape (1)

9. since water freezes at 0°C, so freezing point of the solution containing ethylene glycol will be (1)

since water boils at 100°C, so a solution containing ethylene glycol will boil at 101.024 °C, so it is advisable to

keep this substance in car radiator during summer. (1)

10. (i) The reaction is a zero order reaction.

(ii) For the reaction (½)

(½)

$$-d[A] = k dt$$

integrating both the sides :

$$[A] = kt + C \text{ -----(i)}$$

where C = constant of integration

at t = 0, [A] = [A]₀

Substituting this in equation (i)

$$C = [A]_0$$

Substituting the value of 'C' in equation (i)

$$[A] = -kt + [A]_0$$

$$kt = [A]_0 - [A]$$

$$t = (1)$$

11. The reaction for reducing action of carbon is :

(½)

(½)

$$= + 502 \text{ kJ/mol at } 1273 \text{ K} \text{ (½)}$$

So carbon can be used as reducing agent with MgO(s) at 2273K. (½)

12. The two components of starch are:

(a) Amylose (½)

(b) Amylopectin (½)

Amylose is a straight chain polymer of glucose, while amylopectin is a branched chain polymer of

glucose. (1)

13. (a) On boiling protein of egg gets denatured or coagulated and water of egg get absorbed in it. (1)

(b) Hydrogen bonding between and - NH- groups of peptide bond. (1)

14.

mechanism :

(i) (½)

(ii) (½)

(½)

(iii) (½)

15. (a) (½)

Carbylamine reaction (½)

(b) (½)

Hoffmann bromamide degradation reaction (½)

16. (a) Addition of neutral ferric chloride solution to phenol will give a violet colouration, while no such colouration

will be observed in case of benzyl alcohol. (1)

(b) On addition of Luca's reagent (a mixture of concentrated hydrochloric acid and anhydrous zinc chloride) to

2 - methyl - 2- propanol will give a white turbidity immediately while 2 - Butanol will give turbidity after five minutes. (1)

17. In gaseous phase, basic character of amines increases with increase in number of electron releasing alkyl

groups, due to + I effect, so trend of basic character is (1)

but in aqueous phase, solvation of ammonium cation occurs by water molecules, greater the size of ion, lesser

will be the solvation, and lesser will be the stability of ion, so on combining + I effect and solvation effect, in

aqueous phase trend changes to . (1)

OR

(a) During Friedel-Craft's alkylation, aluminium chloride acts as a catalyst, as well as a Lewis acid, it forms salt

with - NH₂ group of aniline, so that - NH₂ group acquires a positive charge, and acts as a deactivating group,

so aniline does not undergo FCA. (1)

(b) During nitration, in strongly acidic medium aniline is protonated to form anilinium ion, which is a meta directing

group, so along with o- & p- isomers, meta isomer is also obtained. (1)

18. (a) At higher altitudes, partial pressure of oxygen is less than that at ground level, so that oxygen concentration

becomes less in blood or tissues. Hence people suffer from anoxia. (1)

(b) Due to the formation of complex K₂(HgI₄), number of particles in the solution decreases and hence the

freezing point is raised. (1)

19. (½)

Z = 4 (1)

The unit cell is face centered cubic (½)

radius 'r' =

= 141.4 pm. (1)

20. (a) Tetra fluoro ethene (½)

addition polymer (½)

(b) Phenol and formaldehyde (½)

Condensation polymer (1/2)

(c) Isoprene (1/2)

addition polymer (1/2)

21. (a) tetraquadichloro chromium (III) chloride. (1)

(b) $[\text{FeF}_6]_4$ has 4 unpaired electron as F^- is a weak field ligand (1/2)

$[\text{Fe}(\text{CN})_6]_4$ has zero unpaired electron as CN^- is a strong field ligand. (1/2)

(c) Ionisation isomerism. (1/2)

on addition of dilute HCl followed by aqueous will give a white precipitate while the other coordination compound will not give any white precipitate. (1/2)

22. (a) As ferric hydroxide, $\text{Fe}(\text{OH})_3$ is a positively charged sol, so it gets coagulated by chloride ions, Cl^- , released

by NaCl solution. (1)

(b) Cottrell's smoke precipitator, neutraliser the charge on unburnt carbon particles, coming out of chimney and

they get precipitated and settle down at the floor of the chamber. (1)

(c) As physical adsorption, involves only weak vander waal's force of interaction, so many layers of adsorbate

get attached, while chemisorption involves chemical bond formation between adsorbate and adsorbent, so

monolayer is formed. (1)

23. (a) Chlorine water produces nascent oxygen which is responsible for bleaching action and oxidation:

(1)

(b) Both H_3PO_2 and H_3PO_3 have P-H bonds, so they act as reducing agents, but H_3PO_4 , has no P-H bond but

has O-H bonds, so it cannot act as a reducing agent. (1)

(c) Ozone gas acts as a strong oxidising agent, so it oxidises iodide ions to Iodine

I_2 Vapours evolved have violet colour. (1)

24. For first order reaction

(a)

$[\text{R}]_t = 0.23 \text{ M}$ (1)

(1)

(b) (i)

(1)

(ii)

(1/2)

25. (a) 1- Bromo butone, being a primary alkyl halide would react faster by $\text{S}_{\text{N}}2$ pathway, due to less steric

hinderance. (1)

(b) In allyl chloride, the carbocation formed is stabilised due to resonance while the carbocation formed from n-propyl chloride i.e. is less stable, so allyl chloride is more reactive towards nucleophilic substitution reaction. (1)

(c) KCN , being ionic, ions liberated reacts with halo alkanes forming alkyl cyanides but in Ag , being

covalent, does not release ion but lone pair on nitrogen acts as a nucleophile, resulting in formation of

iso cyanides. (1)

26. (a) Nitrogen being smaller in size forms multiple bonding with carbon, So ion is known, but phosphorus does not form bond as it is larger in size. (1)

(b) NO_2 is an odd electron molecule and therefore gets dimerised to stable N_2O_4 . (1)

(c) Because ICl has less bond dissociation enthalpy than I_2

OR

'A' = Sulphur ($\frac{1}{2}$)

B = H_2S gas ($\frac{1}{2}$)

C = SO_2 gas ($\frac{1}{2}$)

D = SO_3 gas ($\frac{1}{2}$)

($\frac{1}{2}$)

($\frac{1}{2}$)

27. (a) Due to antiblood clotting action, aspirin is used for prevention of heart attacks. (1)

(b) As artificial sweeteners provide less calories than natural sweeteners. (1)

(c) Detergents have highly branched hydrocarbon chain, which can not be degraded by bacteria, so they get

accumulated while soap containing straight hydrocarbon chain can be degraded easily (1)

28. (a) As 'A' gives positive iodo form test, so it has ($\frac{1}{2}$)

as 'A' gives positive tollen's test, so it must have – CHO group ($\frac{1}{2}$)

So A is (1)

(b) (i) ($\frac{1}{2}$)

($\frac{1}{2}$)

(ii) ($\frac{1}{2}$)

Acetaldehyde

Butane - 1, 3 -diol ($\frac{1}{2}$)

(iii)

Acetone ($\frac{1}{2}$)

($\frac{1}{2}$)

OR

(a) As 'A' does not give Fehling's or Tollen's test, so it does not have – CHO group but it gives positive iodoform

test and DNP test so it has group (1)

(159)

Q.No. Value Points Marks

So 'A' is :

Acetophenone (1)

B is carboxylic acid obtained by oxidation of A with H_2CrO_4 .

So 'B' is

Benzoic acid (1)

(b) A = (1)

B = ($\frac{1}{2}$)

C = CHI_3 ($\frac{1}{2}$)

29. (a)

(1)

$K_c = \text{antilog}(12.20)$

$= 1.585 \times 10^{12}$ (1)

(b) $M = Z I t$ (1/2)

(x = molar mass of copper)

$x = 63.3 \text{ g/mol}$. (1)

(c) for reaction of tarnished silver ware with aluminium pan is

$(-0.71 \text{ V}) - (-1.66 \text{ V})$ i.e. $+0.95 \text{ V}$ (1)

Tarnished silver ware, therefore, can be cleaned by placing it in an aluminium pan as it is positive. (1/2)

OR

(1(a)) $= (-2.87 \text{ V}) - (1.50 \text{ V})$

$= -4.37 \text{ V}$ (1/2)

$= -6 \times 96500 \times -4.37 \text{ V}$

$= +2350.230 \text{ kJ/mol}$ (1/2)

Since it is positive, reaction is non-spontaneous. (1)

Au^{3+}/Au half cell will be a reducing agent Ca^{2+}/Ca half cell will be an oxidising agent (1/2)

(b) (1/2)

K = specific conductance

$= 40 \text{ Scm}^2 \text{ mol}^{-1}$ (1/2)

(1/2)

$= 1.19 \times 10^{-5}$ (1/2)

30. A = MnO_2 (1/2)

B = K_2MnO_4 (1/2)

C = KMnO_4 (1/2)

(1)

(1/2)

(b) In acidic medium K_2MnO_4 changes to give purple coloured compound along with black precipitate.

Green purple Black (1/2)

compound compound

It is called disproportionation reaction. (1/2)

OR

(a) Due to strong interatomic interaction between unpaired valence electrons. (1)

(b) Because Cl(IV) has extra stability due to empty f_0 orbital (1)

(c) In Mn^{2+} d^5 configuration leads to extra stability of half filled configuration, so $\text{Mn}^{3+}/\text{Mn}^{2+}$ (d^4) tends to get

converted to stable d^5 , configuration of Mn^{2+} , by accepting an electron so $\text{Mn}^{3+}/\text{Mn}^{2+}$ redox couple has more

positive potential than couple (1)

(d) Due to more negative enthalpy of hydration of $\text{Cu}^{2+}(\text{aq})$ than $\text{Cu}^+(\text{aq})$ which compensates for second ionisation

enthalpy of copper. (1)

(e) In the third transition series after lanthanum there is lanthanoid contraction, due to ineffective shielding by

intervening f -orbital electrons and hence second and third transition series elements have similar atomic radii. (1)